

WAKISSHA MOCK-2023 CHEM1 MARKING GUIDE

Name: SSENTAMU FRANK ^{by} Centre/Index No:

School: 0785713136 / 0704131107 Signature: SSENTAMU

SSENTAMU

P525/1
CHEMISTRY
Paper 1
July/August 2023
2 ¼ hours



WAKISSHA JOINT MOCK EXAMINATIONS

Uganda Advanced Certificate of Education

CHEMISTRY

Paper 1

2 hours 45 minutes

Instructions to Candidates

- Attempt all questions in section A and any six questions from section B.
- All questions are to be answered in the spaces provided.
- A Periodic Table with relevant atomic masses is supplied at the end of the paper.
- Mathematical tables (3 figures) and non-programmable silent scientific calculators may be used.
- Illustrate your answers with equations where applicable.
- Molar gas volume at s.t.p = 22.4 dm^3

For Examiner's Use Only

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	Total


© WAKISSHA Joint Mock Examinations 2023

Turn Over

SECTION A (46 MARKS)

Attempt all questions in this section.

1. (a) Sodium -24 which is used as an electrolytic tracer decays by emission of a beta particle and two gamma rays with half-life of 15 hours. (01 mark)
Write the nuclear reaction for the decay of sodium -24.



- (b) 2.4g of sodium -24 were allowed to disintegrate for 72 hours. Calculate the mass of the radioactive isotope that decayed. (04 marks)

From $t_{1/2} = \frac{0.693}{k}$ From $2.303 \log\left(\frac{N_0}{N_t}\right) = kt$

$k = \frac{0.693}{15}$ $2.303 \log\left(\frac{2.4}{N_t}\right) = 0.0462 \times 72$

$k = 0.0462$ $\log\left(\frac{2.4}{N_t}\right) = \frac{0.0462 \times 72}{2.303}$

$\frac{2.4}{N_t} = 10^{1.4443764}$ $N_t = 0.0863$

Amount decayed = $2.4 - 0.0863 = 2.3137\text{g}$

- (c) State any two other uses of radioactive isotopes.


- Radioactive oxygen-18 is used in determination of mechanism of esterification reaction.

- Radioactive carbon-14 is used in carbon dating.


2. The atomic number of chromium is 24.

- (a) Write the;

- (i) electronic configuration of chromium. (01 mark)




- (ii) formulae of three common oxides of chromium. (1½ marks)




- (b) The oxide(s) in a(ii) are either basic, amphoteric or acidic. Write an equation for the reaction between the;


- (i) basic oxide of chromium and dilute mineral acids. (1½ marks)



- (ii) acidic oxide of chromium and sodium hydroxide solution. (1½ marks)




3. Complete the following equations and in each case outline a suitable mechanism for the reaction.




Mechanism:



(Accept formation of ethene and mark correspondingly. E2 mechanism)



Mechanism:




4. (a) Define the term common ion effect. (01 mark)

Is the precipitation of a salt from its saturated solution by adding an aqueous solution of a soluble salt containing a common ion to the ions in the equilibrium of the sparingly soluble salt.

(b) The solubility of Lead (II) chloride in 0.02 M calcium chloride at 25°C is $2.951625 \text{ gdm}^{-3}$.

(i) Calculate the solubility of lead (II) chloride in gdm^{-3} in pure water at 25°C. (05 marks)

RFM of PbCl_2
 $= 207 + 35.5 \times 2$
 $= 278$



$[\text{PbCl}_2] = \frac{2.951625}{278}$
 $= 0.106 \text{ M}$

$K_{sp} = [\text{Pb}^{2+}][\text{Cl}^-]^2$
 $= 0.106 \times (0.04)^2$
 $K_{sp} = 1.696 \times 10^{-4} \text{ mol}^3 \text{dm}^{-3}$

From $\text{PbCl}_2(\text{s}) \rightleftharpoons \text{Pb}^{2+}(\text{aq}) + 2\text{Cl}^-(\text{aq})$ Let the solubility in water be $x \text{ mol dm}^{-3}$.

$K_{sp} = [\text{Pb}^{2+}][\text{Cl}^-]^2$

$K_{sp} = x(2x)^2$
 $K_{sp} = 4x^3$
 $x = \sqrt[3]{\frac{1.696 \times 10^{-4}}{4}} = \sqrt[3]{4.24 \times 10^{-5}}$
 $x = 3.487 \times 10^{-2}$ (1/2 mark)

(ii) Comment on your answer in b(i) above.

Lead (II) chloride has a higher solubility in water than in 0.02M calcium chloride. 0 1/2

RFM of $\text{PbCl}_2 = 278$
 Turn Over
 1 mole weighs 278g
 $3.487 \times 10^{-2} \text{ moles weigh } 278 \times 3.487 \times 10^{-2}$
 $= 12.15 \text{ gdm}^{-3}$

CS CamScanner

$= 9.69386 \text{ gdm}^{-3}$


5. The enthalpies of some chemical reactions are given below.

Reaction	ΔH° (KJmol ⁻¹)
(i) $\text{MgO}_{(s)} + 2\text{HCl}_{(aq)} \longrightarrow \text{MgCl}_{2(aq)} + \text{H}_2\text{O}_{(l)}$	-146.2
(ii) $\text{Mg}_{(s)} + 2\text{HCl}_{(aq)} \longrightarrow \text{MgCl}_{2(aq)} + \text{H}_2_{(g)}$	-478.4
(iii) $2\text{H}_{2(g)} + \text{O}_{2(g)} \longrightarrow 2\text{H}_2\text{O}_{(l)}$	-572

(03 marks)

(a) Calculate the enthalpy of formation of magnesium oxide.

Required equation



$\Rightarrow \text{Mg}_{(s)} + \frac{1}{2}\text{O}_{2(g)} \longrightarrow \text{MgO}_{(s)} \quad \Delta H = -478.4 + \frac{1}{2}(-572) + 146.2$

$\Delta H = -618.2 \text{ KJmol}^{-1}$

(b) State whether Magnesium oxide is stable or not. Give a reason for your answer. (1 1/2 marks)

Magnesium oxide is stable since it has a 1/2 negative enthalpy of formation.

6. The physical properties of the hydrides of fluorine and Iodine are shown below.

Hydride	HF	HI
Boiling point	+19.9°C	-35.1°C
Physical state	Liquid	Gas

(a) Explain the variation in physical properties of the hydrides. (02 marks)

Hydrogen fluoride is a liquid at room temperature because due to the high electronegativity of fluorine the molecules of hydrogen fluoride are held together by strong hydrogen bonds which can't be overcome at room temperature hence high boiling point and liquid state. In hydrogen iodide, iodine has very low electronegativity hence its molecules are held together by weak van der Waal forces which are easily overcome hence existing as a gas with very low boiling point.

(b) Describe the reactions of the hydrides with:

(i) sodium carbonate solution. (1 1/2 marks)

The hydrides react with sodium carbonate solution to form salts, water and carbonate gas. For example,

$\text{Na}_2\text{CO}_{3(aq)} + 2\text{HI}_{(aq)} \longrightarrow 2\text{NaI}_{(aq)} + \text{H}_2\text{O}_{(l)} + \text{CO}_{2(g)}$

Hydrogen fluoride however forms a salt of sodium hydrogen bifluoride (NaHF_2).


(ii) concentrated sulphuric acid.

(1 1/2 marks)

There is no reaction between hydrogen fluoride and concentrated sulphuric acid, but concentrated sulphuric acid oxidises hydrogen to diiodine and itself reduced to sulphur dioxide.
 $H_2SO_4(l) + 2HI(g) \rightarrow 2H_2O(l) + SO_2(g) + I_2(s)$
(Accept reduction to H_2S , S , I_2 and H_2O).

7. Methylbenzoate was warmed with excess acidified water to form two organic products which were separated by distillation.

(a) Write an equation for the reaction that took place. (01 mark)




(b) Name a reagent that can be used to distinguish the products in (a) above. State what would be observed if each of the products is separately treated with the reagent you have named. (03 marks)

Reagent.

Sodium carbonate solution ✓

Observation.



Bubbles of a colourless gas that turns lime water milky ✓

CO: No observable change ✓


(c) State one other physical method by which the products of the reaction can be separated. (1/2 mark)

By filtration ✓

8. Compound E expands normally and has a critical temperature of $31^\circ C$ at 73.9 atm pressure. The triple point of E is $-57^\circ C$ at 5.2 atm pressure.

(03 marks)

(a) Sketch a well-labelled phase diagram of E.



Turn Over
5

(b) State what would happen when I; at; (01 mark)

(i) 180 K temperature and 50 atm pressure was heated at constant pressure. (01 mark)

When heated at constant pressure
it will melt to liquid state. ✓ 01

(ii) -57°C and 5.2 atm pressure was compressed at constant temperature. (01 mark)

When compressed at constant temperature
E, will solidify. ✓ 01

9. Lead (IV) oxide reacts with excess ice cold concentrated hydrochloric acid to form a complex liquid which forms a yellow precipitate on addition of a saturated solution of ammonium chloride. The dry precipitate reacts with concentrated sulphuric acid to form a pale yellow liquid.

(a) Name the;

(i) complex liquid. (½ mark)

hexachloroplumbic(IV) acid ✓ ½


(ii) yellow precipitate. (½ mark)

Ammonium hexachloroplumbate(IV) ✓ ½

(iii) pale yellow liquid. (½ mark)

Lead(IV) chloride ✓ ½

(b) Write an equation for the reaction between water and the pale yellow liquid (01 mark)









(c) Name the type of reaction that occurs in (b). (½ mark)

Hydrolysis ✓ ½


SECTION B (54 MARKS)

Attempt any six questions from this section.

10. Write equations to show how the following conversions can be effected.



11. (a) 0.05M copper (II) sulphate was titrated with aqueous ammonia. The conductivity of the mixture varies as shown by the graph below.



Turn Over

State the reason(s) why;

- (i) conductivity is initially high at point A. (01 mark)


Copper (II) sulphate is a strong electrolyte which ionises completely to give many copper (II) ions and sulphate ions hence high conductivity.

- (ii) conductivity almost remains constant along CD. (01 mark)


The ionisation of weak ammonia solution is suppressed by presence of a common ion (hydroxide ion) shifting the equilibrium $\text{NH}_3(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightleftharpoons \text{NH}_4^+ + \text{OH}^-$ to the left.

- (b) Write an equation for the reaction that takes place along;

- (i) AB (01 mark)



- (ii) BC (01 mark)



- (c) The electrolytic conductivity of water at 25°C is $5.484 \times 10^{-8} \Omega^{-1} \text{cm}^{-1}$ and its concentration is 18 g per 18 cm³. Given that the molar conductivity at infinite dilution of H⁺ and OH⁻ are 349.8 and 198.6 $\Omega^{-1} \text{cm}^2 \text{mol}^{-1}$ respectively. Calculate the;

- (i) degree of ionization of water at 25°C. (3½ marks)

$$\begin{aligned} \Lambda_0 \text{H}_2\text{O} &= \Lambda_0 \text{H}^+ + \Lambda_0 \text{OH}^- \\ &= 349.8 + 198.6 \\ &= 548.4 \Omega^{-1} \text{cm}^2 \text{mol}^{-1} \end{aligned}$$

$$\Lambda_c = \frac{1000 \times 5.484 \times 10^{-8}}{55.56}$$

$$\Lambda_c = \frac{5.484 \times 10^{-5}}{55.56}$$

$$\Lambda_c = 9.87 \times 10^{-7} \Omega^{-1} \text{cm}^2 \text{mol}^{-1}$$

$$\alpha = \frac{\Lambda_c}{\Lambda_0} = \frac{9.87 \times 10^{-7}}{548.4}$$

$$\alpha = \frac{1.8 \times 10^{-9}}{55.56}$$


- (ii) ionic product of water, Kw at 25°C. (1½ marks)

$$K_w = C^2 \alpha^2 \text{ where } \alpha \text{ is degree of ionisation of water.}$$

$$K_w = 55.56^2 (1.8 \times 10^{-9})^2$$

$$K_w = 1.0 \times 10^{-14} \text{ mol}^2 \text{dm}^{-6}$$

12. Dilute nitric acid reacts with phenol according to the equation;



The products were separated by steam distillation.

(a) State the reason(s) why the:

(i) reaction occurs with dilute nitric acid in the absence of a catalyst unlike with benzene. (02 marks)

In phenol unlike benzene; presence of a hydroxyl group activates the benzene ring due to increased number of delocalised electrons making it more susceptible to electrophilic attack hence reaction with nitric acid in absence of a catalyst.


(ii) two products can be separated by steam distillation. (02 marks)

It's because 4-nitrophenol is non-volatile (has a very high boiling point) owing to the strong hydrogen bonds which can't easily be broken hence it can be separated from the volatile 2-nitrophenol which is also immiscible with water by steam distillation.

(b) When the mixture was steam distilled at 1.0 atm at 96°C, the mass of water in the steam distillate was 0.90 g. Calculate the mass of the second component of the distillate.

(Saturated vapour pressure of water at 96°C = 0.825 atm). (03 marks)

Second component is 2-nitrophenol



$$P_x = 1.0 - 0.825 = 0.175 \text{ atm}$$

$$\frac{M_{\text{H}_2\text{O}}}{M_x} = \frac{P_{\text{H}_2\text{O}} \cdot M_{\text{H}_2\text{O}}}{P_x \cdot M_x}$$

$$\frac{0.90}{M_x} = \frac{0.825 \times 18}{0.175 \times 139}$$

$$M_x = \frac{0.90 \times 0.175 \times 139}{0.825 \times 18}$$

$$M_x = 1.474 \text{ g}$$

(c) State two advantages of steam distillation. (02 marks)

- It is useful in purifying organic compounds which tend to decompose near their boiling points.
- It prevents thermal decomposition of compounds which occurs at high temperature.

Turn Over

13. (a) Name one reagent that can be used to distinguish between the following pairs of compounds. State what would be observed when each member of the pair is separately treated with the reagent you have named.

(i) $K_2SO_4(aq)$ and $K_3PO_4(aq)$ (02 marks)

Reagent

Lead (II) nitrate solution and dilute nitric acid

Observation

K_2SO_4 : white precipitate insoluble in acid ✓ 02

K_3PO_4 : white precipitate soluble in acid ✓

(ii) $NaCl(aq)$ and $Na_2C_2O_4(aq)$ (02 marks)

Reagent

Calcium chloride solution ✓

Observation

$NaCl$: No observable change ✓ 02


$Na_2C_2O_4$: white precipitate ✓

(b) Explain each of the following observations.

(i) When sodium hydroxide solution is added to neutral potassium dichromate solution, the orange solution turns yellow and pale yellow precipitate is formed on addition of lead (II) nitrate solution.


In presence of sodium hydroxide solution, potassium dichromate solution is converted to potassium chromate (VI) solution which is yellow. 2½ marks

On addition of lead (II) nitrate solution, lead chromate (VI) is formed which is a yellow precipitate. 2½



(ii) Manganese (II) sulphate solution in the presence of concentrated nitric acid forms a purple solution on addition of sodium bismuthate.

(2½ marks)



Bismuthate (V) ions oxidise manganese (II) ions in manganese (II) sulphate solution to manganate (VII) ions which are purple in colour. 2½

09


14. An organic compound Q has a molecular formula; $C_4H_8O_2$ Corrected from C_4H_8O to $C_4H_8O_2$

Q has the following chemical properties;

- forms a yellow precipitate with both 2,4-dinitrophenyl hydrazine and aqueous iodine in the presence of sodium hydroxide.
- forms a cloudy solution after 8 minutes on addition of a solution of anhydrous zinc chloride in concentrated hydrochloric acid.
- gives a silver mirror on addition of ammoniacal silver nitrate solution.

(a) Write the;

(i) structural formula of Q. OH ✓ (01 mark)




(ii) IUPAC name of Q. (01 mark)


3-hydroxybutanal ✓

(b) Write an equation for the reaction between Q and;


(i) anhydrous zinc chloride in the presence of concentrated hydrochloric acid. OH Cl (01 mark) ✓




(ii) ammoniacal silver nitrate solution. OH (01 mark)





(iii) saturated sodium hydrogensulphite solution OH (01 mark)



(c) Suggest a suitable mechanism for the reaction between Q and acidified 2,4-dinitrophenyl hydrazine. OH (04 marks) ✓



Let $\text{2,4-dinitrophenyl} \text{ be } P$.



Turn Over
CS CamScanner
09

15. (a) Ammonia is obtained on large scale in the Haber process according to the equation.



State the effect of the following on the yield of ammonia. Give a reason for your answer.

- (i) high pressure of 150 - 200 atm. (1½ marks)

The forward reaction occurs with a decrease in volume (moles) hence favoured by high pressure which shifts equilibrium position to the right to form more ammonia increasing its yield.


- (ii) high temperature above 450°C (1½ marks)

The forward reaction is exothermic and hence favoured by low temperature. Increasing temperature above 450°C favours the backward reaction hence decreasing the yield of ammonia.


- (b) 3.0 moles of nitrogen gas were mixed with 1.0 moles of hydrogen gas in a 500 cm³ bulb. The mixture was allowed to attain equilibrium at 450°C and the mass of ammonia in the equilibrium mixture was found to be 0.34 g. Calculate the concentration equilibrium constant, K_c at 450°C. (03 marks)

Moles of NH₃ = $\frac{0.34}{17} = 0.02$ moles
 Moles of N₂ = $3 - 3(0.01) = 2.97$ moles
 H₂ = $1 - 0.01 = 0.99$ moles

Initial moles
 If x moles react
 moles at equilibrium



- (c) Write equations to show how the ammonia obtained from the Haber process can be converted to nitric acid. (03 marks)



16. (a) Lithium belongs to group I of the Periodic Table but its properties resemble Magnesium of group II. State three:

- (i) reasons why the chemistry of Lithium differs from other group I elements. (1½ marks)


- Has very small atomic radius
- Has very high electropositivity
- Lithium ion has very high polarizing power
- Lithium has very low electronegativity

(ii) properties of lithium to show how its chemistry resembles that of Magnesium. (03 marks)


- Both Lithium and magnesium react with oxygen in air to form normal oxides ✓
- Both form carbonates that decompose on heating ✓
- Both form nitrates that decompose on heating to form normal oxides ✓

(b) Write an equation for the reaction between;

(i) Magnesium nitride and water. (1½ marks) ✓



(ii) Beryllium chloride and sodium hydroxide solution. (1½ marks) ✓



(iii) Barium peroxide and dilute hydrochloric acid. (1½ marks) ✓



09

17. (a) Ethanol and hexane form an azeotropic mixture of composition 38.42% ethanol and 61.58% hexane. The density of the azeotrope is 0.687 g cm^{-3} .

Substance	Ethanol	Hexane	Azeotrope
Boiling point (°C)	78.4	68.9	59.15

(i) State the type of deviation from Raoult's law. (01 mark)

Positive deviation ✓ 01


(ii) Explain your answer in (a)(i) above. (02 marks)

The mixture forms an azeotrope with minimum boiling point hence it contributes more vapour than expected from Raoult's law due to very weak forces of attraction between the components hence positive deviation. ✓ 02

Turn Over

13

(b) (i) Sketch a well-labelled boiling point composition diagram for the ethanol-hexane system. (02 marks)



(ii) A mixture containing 25% liquid ethanol was fractionally distilled. Identify the substance obtained as;

- distillate... Azeotrope $\frac{1}{2}$ (1/2 mark)
- residual liquid... Pure hexane $\frac{1}{2}$ (1/2 mark)

(c) 50 cm³ of the azeotrope was shaken with 100 cm³ of choline chloride (solvent) at 25°C. Calculate the mass of ethanol extracted by choline solvent. (Partition coefficient of ethanol between choline chloride and hexane at 25°C is 15.80). (03 marks)

Amount of ethanol

1 cm³ of azeotrope contains 0.687g
 50 cm³ of azeotrope contains (0.687 x 50)g
 = 34.35g

% ethanol = $\frac{34.35}{100} \times 100 = 34.35\%$

K.D = $\frac{[\text{ethanol}]_{\text{choline chloride}}}{[\text{ethanol}]_{\text{hexane}}}$

Let the mass of ethanol extracted by choline chloride be x .

$$15.80 = \frac{x}{\frac{13.197 - x}{50}}$$

$$15.80 = \frac{x}{100} \times \frac{50}{13.197 - x}$$

$$15.80 = \frac{x}{2(13.197 - x)}$$

$$417.0252 - 31.6x = x$$

$$\frac{417.0252}{32.6} = \frac{32.6x}{32.6}$$

$$x = 12.792\text{g of ethanol}$$

09

THE PERIODIC TABLE

1	2											3	4	5	6	7	8	
I H 1.0																I II 1.0	2 He 4.0	
3 Li 6.9	4 Be 9.0											5 B 10.8	6 C 12.0	7 N 14.0	8 O 16.0	9 F 19.0	10 Ne 20.2	
11 Na 23.0	12 Mg 24.3											13 Al 27.0	14 Si 28.1	15 P 31.0	16 S 32.1	17 Cl 35.4	18 Ar 40.0	
19 K 39.1	20 Ca 40.1	21 Sc 45.0	22 Ti 47.9	23 V 50.9	24 Cr 52.0	25 Mn 54.9	26 Fe 55.8	27 Co 58.9	28 Ni 58.7	29 Cu 63.5	30 Zn 65.7	31 Ga 69.7	32 Ge 72.6	33 As 74.9	34 Se 79.0	35 Br 79.9	36 Kr 83.8	
37 Rb 85.5	38 Sr 87.6	39 Y 88.9	40 Zr 91.2	41 Nb 92.9	42 Mo 95.9	43 Tc 98.9	44 Ru 101	45 Rh 103	46 Pd 106	47 Ag 108	48 Cd 112	49 In 115	50 Sn 119	51 Sb 122	52 Te 128	53 I 127	54 Xe 131	
55 Cs 133	56 Ba 137	57 La 139	72 Hf 178	73 Ta 181	74 W 184	75 Re 186	76 Os 190	77 Ir 192	78 Pt 195	79 Au 197	80 Hg 201	81 Tl 204	82 Pb 207	83 Bi 209	84 Po (209)	85 At (210)	86 Rn (222)	
87 Fr (223)	88 Ra (226)	89 Ac (227)																
			59 La 139	60 Ce 140	61 Pr 141	62 Nd 144	63 Pm (145)	64 Sm 152	65 Eu 152	66 Gd 157	67 Tb 159	68 Dy 162	69 Ho 165	70 Er 167	71 Tm 169	72 Yb 173	73 Lu 175	
			89 Ac (227)	90 Th 232	91 Pa 231	92 U 238	93 Np 237	94 Pu (244)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf 251	99 Es (254)	100 Fm (257)	101 Mv (256)	102 No (254)	103 Lw	

1. I → Indicates atomic number.
II
2. II
1.0 ← Indicates relative atomic mass.

END