

UNEB UACE APPLIED MATHS 2018

13. a) The first three terms of a Geometric Progression (G.P) are 4,8 and 16. Determine the sum of the first ten terms of the G.P
 b) An arithmetic Progression (A.P) has a common difference of 3.A Geometric Progression (G.P) has a common ratio of 2. A sequence is formed by subtracting the terms of the A.P from the corresponding terms of the G.P. The third term of the sequence is 4. The sixth term of the sequence is 79. Find the first term of the
 i) A.P
 ii) G.P

14. Evaluate:

a) $\int_0^{\frac{\pi}{2}} \sin 5x \cos 3x \, dx$

b) $\int_0^{\sqrt{5}} \frac{dx}{9+4x^2}$

15. The line $y= mx + c$ is a tangent to the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

- a) Obtain an expression for c in terms of a, b, and m
- b) Calculate the gradients of the tangents to the ellipse through the

$$\left(\sqrt{(a^2 + b^2)}, 0\right)$$

16. The rate at which the temperature of a body falls is proportional to the difference between the temperature of the body and that of its surrounding. The temperature of the body is initially 60°C. After 15minutes the temperature of the body is 50°C. The temperature of the surrounding is 10°C.
 a) Form a differential equation for the temperature of the body.
 b) Determine the time it takes for the temperature of the body to reach 30°C.

SECTION A

1. A stone is thrown vertically upwards with a velocity of 21ms⁻¹
 Calculate the:
 a) maximum height attained by the stone.
 b) time the stone takes to reach the maximum height.

2. Two events A and B are such that

$$P(A/B) = 2/5, P(B) = 1/4 \text{ and } P(A) = 1/5$$

Find:

- a) P(A#B)
- b) P(A#B)

3. The table below shows how T varies with S

T	-2.9	-0.1	2.9	3.1
S	30	20	12	9

Use linear interpolation/ extrapolation to estimate the value of

- a) T when S = 26
- b) S when T =3.4

4. A particle of mass 15kg is pulled up a smooth slope by a light inextensible string parallel to the slope. The slope is 10.5m long and inclined at $\sin^{-1}(\frac{4}{7})$ to the horizontal. The acceleration of the particle is 0.98ms⁻².
 Determine the:
 a) tension in the string
 b) work done against gravity when the particle reaches the end of the slope.

5. The price index of an article in 2000 based on 1998 was 130. The price index for the article in 2005 based on 2000 was 80.
 Calculate the:

- a) price index of the article in 2005 based on 1998.
- b) price of the article in 1998 if the price of the article was 45,000 in 2005.

6. Two numbers A and B have maximum possible errors e_a and e_b respectively.

- a) Write an expression for the maximum possible error in their sum
- b) If A = 2.03 and B = 1.547, find the maximum possible error in A+B.

7. In an equilateral triangle PQR, three forces of magnitude 5N, 10N and 8N act along the sides PQ, QR and PR respectively. Their directions are in the order of the letters. Find the magnitude of the resultant force.

8. A biased coin is such that a head is three times as likely to occur as a tail. The coin is tossed 5 times. Find the probability that at most two tails occur.

SECTION B

Answer any five questions from this section.

9. The frequency distribution below shows the ages of 240 students admitted to a certain University.

Age (years)	Number of Students
18 - <19	24
19 - <20	70
20 - <24	76
24 - <26	48
26 - <30	16
30 - <32	6

- a) Calculate the mean age of the students.
- b) i) Draw a histogram for the given data
- ii) Use the histogram to estimate the modal age.

10. A particle of mass 4kg starts from rest at a point $(2\mathbf{i} - 3\mathbf{j} + \mathbf{k}) \text{ m}$. It moves with acceleration $\mathbf{a} = (4\mathbf{i} + 2\mathbf{j} - 3\mathbf{k})\text{ms}^{-2}$ when a constant force \mathbf{F} acts on it.

Find the:

- a) force \mathbf{F}
- b) velocity at any time t
- c) work done by the force \mathbf{F} after 6 seconds.

11. a) Use the trapezium rule with 6 – ordinates to estimate the value of correct to three decimal places.

b) i) Evaluate

$$\int_0^{\frac{\pi}{2}} (x + \sin x) dx$$

correct to three decimal places

ii) Calculate the error in your estimation in (a) above.


$$\int_0^{\frac{\pi}{2}} (x + \sin x) dx$$

iii) Suggest how the error may be reduced.

12. A random variable X has a normal distribution where $P(X > 9) = 0.9192$ $P(X > 11) = 0.7580$. Find:

- a) the values of the mean and standard deviation
- b) $P(X > 10)$

13. The figure below shows a uniform beam of length 0.8 meters and mass 1 kg. The beam is hinged at A and has a load of mass 2kg attached at B.



The beam is held in a horizontal position by a light inextensible string of length 0.5 meters. The string joins the mid-point M of the beam to a point C vertically above A.

Find the :

- a) tension in the string.
- b) magnitude and direction of the force exerted by the hinge.

14. a) Draw on the same axes the graphs of the curves $y = 2 - e^{-x}$ and $y = \sqrt{x}$ for $2 \leq x \leq 5$.

b) Determine from your graphs the interval within which the root of the equation $e^{-x} + \sqrt{x-2} = 0$ lies. Hence, use Newton – Raphson’s method to find the root of the equation correct to 3 decimal places.

15. The table below shows the number of red and green balls put in three identical boxes A, B and C.

Boxes	A	B	C
Red balls	4	6	3
Green balls	2	7	5

A box is chosen at random and two balls are then drawn from it successively without replacement. If the random variable X is “the number of green balls drawn”,

- a) draw a probability distribution table for X
- b) calculate the mean and variance of X

16. At 10:00am, ship A and ship B are 16km apart. Ship A is on a bearing $N35^{\circ}E$ from ship B. Ship A is travelling at 14kmh^{-1} on a bearing $S29^{\circ}E$. Ship B is travelling at 17kmh^{-1} on a bearing $N50^{\circ}E$. Determine the;

- velocity of ship B relative to ship A
- closest distance between the two ships and the time when it occurs.

PURE MATHEMATICS – PAPER 1

1. Given that

$$Q = \sqrt{80 - 0.1P} \text{ and } E = \frac{-dQ}{dP} \cdot \frac{P}{Q}, \text{ find } E \text{ when } P = 600$$

2. Find the area enclosed between the curve $y = 2x^2 - 4x$ and the xx -axis

3. Solve $2\cos 2\theta - 5\cos \theta = 4$ for $0^{\circ} \leq \theta < 360^{\circ}$

4. Given that $\alpha + \beta = \frac{-1}{3}$ and $\alpha\beta = \frac{2}{3}$, form a quadratic equation whose roots are $\frac{\alpha}{\beta}$ and $\frac{\beta}{\alpha}$

5. Given the plane $4x + 3y - 3z - 4 = 0$;

a) show that the point A (1,1,1) lies on the plane.

b) find the perpendicular distance from the plane to the point B (1,5,1)

6. Find the equation of the tangent to the curve

$$y = \frac{a^3}{x^2} \text{ at the point } P \left(\frac{a}{t}, at^2 \right)$$

7. Find the area enclosed between the curve $y = 2x^2 - 4x$ and the x -axis

8. Show that the modulus of

$$\frac{(1-i)^6}{1+i} = 4\sqrt{2}$$

9. a) Determine the perpendicular distance of the point (4,6) from the line $2x + 4y - 3 = 0$

b) Show that the angle θ , between two lines with gradients m_1 and m_2 is given by

$$\theta = \tan^{-1} \left(\frac{m_1 - m_2}{1 + m_1 m_2} \right)$$

Hence find the acute angle between the lines