

Student's Name:

Signature:.....

BIOLOGY
PAPER 1
(Theory)
3 Hours



KANGULUMIRA PUBLIC SECONDARY SCHOOL
SENIOR SIX
TRIAL TEST 003

BIOLOGY
Paper 1
(Theory)
3 HOURS

INSTRUCTIONS TO CANDIDATES:

*This paper consists of **two** sections: **A** and **B**. it has **SIX** examination items.*

*Section **A** has **Two Compulsory** items.*

*Section **B** has **Four ITEMS**: Answer two **ITEMS**, choosing one item from each part.*

*Answers to section **A** **must** be written in the spaces provided while answers to **section B** must be written in the answer booklet (s) provided.*

*Answer **FOUR** items in all.*

*Any additional item(s) answered will **not** be scored.*

SECTION A

Attempt all the items in this section

ITEM 1

A rural community started a small cassava flour processing factory to improve food security. After six months, workers began complaining of headaches, dizziness, and muscle weakness. Some fish in a nearby stream were also found dead. The factory uses crushed cassava soaked in water before drying. During processing, a bitter Smell similar to almonds was noticed. The wastewater from the factory is discharged into the stream.

Further laboratory tests produced the following results:

Parameter	Normal fish	Fish near factory
Dissolved oxygen (mg/L)	9.0	3.5
ATP per mitochondrion	28	12
Cytochrome C oxidase activity (%)	100	30
Rate of glycolysis	Normal	Increased
Mutation rate in liver cells (%)	2	18

Additional findings:

- Cyanide was detected in wastewater.
- High temperature recorded in discharged water.
- Fish gill epithelial cells showed damaged plasma membranes.
- Ribosomes in liver cells appeared detached from rough ER.

Task:

- (a) Analyze how factory wastewater disrupts fish cell function by affecting mitochondrial structure and ATP production, inhibiting enzymes, damaging plasma membranes, and increasing DNA mutation rates through impaired protein synthesis.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

ITEM 2:

Tea farmers in Kamuli pruned their plants heavily just before the cold season to increase branching. However, after pruning, plants showed slow recovery, smaller leaves, and delayed flowering.

Kamuli experiences cool temperatures (14-16⁰C), Frequent fog, and shorter daylight hours during this period.

Data collected:

Parameter	Before Pruning	After Pruning (Cold Season)
Leaf area	Large	Small
CO ₂ uptake	Moderate	Low
Chloroplast activity	Normal	Reduced
Gibberellin level	Moderate	Low
Cytokinin level	Moderate	Low
Growth rate	Moderate	Very slow

Task:

- (a) Analyze how low temperature combined with pruning affects photosynthesis, hormone balance, primary growth, and photoperiod response.

.....

.....

.....

.....

For responses Whatsapp: 0788060585 @ Ugx 3000

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Evaluate how pruning and seasonal management should be adjusted to improve yield.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SECTION B

Part I

Attempt one item from this section.

ITEM 3:

At a secondary school in Mbale (Altitude 1,200m), students participated in a tree-planting exercise near a busy roadside. After several hours, four students showed unusual signs.

- Musa felt dizzy and had a headache.
- Lillian developed itchy eyes and sneezing.
- Tom reacted slowly when called and staggered slightly.
- Aisha complained of blurred vision in dim light.

The health club recorded the following:

Parameter	Musa	Lillian	Tom	Aisha	Normal
Blood oxygen saturation (%)	82	96	88	95	95-100
Carboxyhaemoglobin (%)	18	1	10	1	<2
Heart rate (beats/ min)	120	90	110	85	70-100
Reaction time (s)	0.32	0.19	0.30	0.18	0.18
Tear secretion	Normal	High	Normal	Normal	Normal
Rod cell response (retina test)	Normal	Normal	Normal	Low	Normal

Later it was discovered that traffic had been heavy and some students had not eaten or drunk water all morning.

Task:

- (a) Using the data, analyze how carbon monoxide exposure, dehydration, allergic reactions, and sensory receptor function affected the students' oxygen transport, heart activity, nervous coordination, and vision.
- (b) Evaluate practical measures to prevent such conditions during outdoor school activities.

ITEM 4:

During a school mountain hiking trip to the slopes of Mount Elgon (elevation 3000m), students participated in a survival challenge. The weather was cold (10°C), and the air was thin.

Two students, Musa and Joel, showed unusual responses:

- Musa complained of headache and dizziness. His lips appeared slightly blue.
- Joel fainted briefly near a charcoal stove used for cooking inside a tent.
- Both had increased breathing rates.
- Later that night, some students developed itchy rashes after eating packed groundnuts.
- One student, Aisha, had recently received a tetanus booster vaccine.
- Rapid malaria test kits were used on two febrile students and were negative.

The teacher recorded the following:

Parameter	Musa	Joel
Breathing rate (breaths/min)	30	28
Pulse rate (beats/min)	120	130
Blood oxygen saturation (%)	82	78
Carboxyhaemoglobin level (%)	2	25
Core body temperature ($^{\circ}\text{C}$)	35.5	36.8
Urine output	Low	Normal

The school nurse suspected problems related to oxygen transport, temperature regulation, immunity, and nerve function.

Task:

- (a) Analyze the physiological causes of Musa's and Joel's symptoms by linking oxygen dissociation, haemoglobin function, carbonmonoxide effects, nervous and hormonal control of heart rate, hypothalamic temperature regulation, histamine action, vaccination, and rapid test kit mechanisms.
- (b) Propose and evaluate preventive and management strategies for future mountain hikes.

PART II

Attempt one item from this section.

ITEM 6:

In the bean-growing regions of eastern Uganda, farmers are battling an outbreak of black bean aphids (*Aphis fabae*). A researcher from Makerere University found that the aphids exist in two main strains: One that is resistant to a common pyrethroid pesticide, and one that is susceptible. The resistance is controlled by a dominant allele (R). The aphids also vary in their colour: some are shiny black, while others are dull brown. The colour is controlled by a separate gene (B for black, dominant over b for brown). The researcher performed a cross between a true-breeding resistant, black aphid and a true-breeding susceptible, brown aphid. The F1 generation were all resistant and black. However, when these F1 males were crossed with susceptible, brown females (a test cross), the offspring were not in a 1:1:1:1 ratio. The 500 offspring produced were:

- Resistant, Black: 205
- Resistant, Brown: 45
- Susceptible, Black: 48
- Susceptible, Brown: 202

The researcher also noted that ladybird beetles (a natural predator of aphids) prefer to eat the brown aphids, as they are more visible the green bean leaves.

Task:

- (a) Analyze the genetic data from the test cross to explain the inheritance pattern of the resistance and colour genes, predict how the interaction between pesticide use and predator preference will affect the allele and phenotype frequencies in the aphid population over time.
- (b) Propose and evaluate integrated pest management strategies for farmers that combines chemical and biological controls, sustainably.

ITEM 7:

In the rice-growing region of Kamuli district, farmers have noticed that brown planthoppers are causing reduced yields despite pesticide application. Observations include:

- Some planthoppers carry alleles for pesticide resistances showing incomplete dominance.
- Resistant allele frequency increased from 10% to 55% over four years.
- Paddy fields with high bird populations report fewer planthoppers, while frog populations have declined.

For responses Whatsapp: 0788060585 @ Ugx 3000

- Waterlogged fields favour planthopper nymph survival, increasing over the wet season.
- Some planters practice crop rotation; others leave paddies fallow.

Task:

- Analyze the genetic, ecological, and population factors contributing to the increase in pesticide-resistant planthoppers.
- Propose and evaluate sustainable, biologically informed pest management strategy for Kamuli rice fields.

STAY BLESSED MY PEOPLE

For responses Whatsapp: 0788060585 @ Ugx 3000



WHAT WE DO?

1. Website development
2. Logo design
3. Social media graphics
4. Printing & branding
5. Banner, Flyers & posters
6. Profile / portfolio design
7. Software dev't
8. Book layouts
9. Business registrations
10. Proposals
11. Professional CV's & Resumes

AFFORDABLE PRICES

Get Discount On Bulky Orders

FOR MORE INFORMATION MESSAGE US:

0788060585



EXODUS PUBLISHERS